



YRBSS

National Youth Risk Behavior Survey

Trends in the Prevalence of Tobacco Use

What Is the National Youth Risk Behavior Survey (YRBS)?

The national YRBS monitors priority health risk behaviors that contribute to the leading causes of death, disability, and social problems among youth and adults in the United States. The national YRBS is conducted every two years during the spring semester and provides data representative of 9th through 12th grade students in public and private schools throughout the United States.

1991	1993	1995	1997	1999	2001	Changes from 1991 – 2001 ¹	Change from 1999 – 2001 ²
Current smokeless tobacco use (Used chewing tobacco, snuff, or dip on one or more of the 30 days preceding the survey.)							
NA ³	NA	11.4 (±1.7) ⁴	9.3 (±2.2)	7.8 (±2.2)	8.2 (±1.5)	Decreased, 1995 – 2001	No change
Smokeless tobacco use on school property (Used chewing tobacco, snuff, or dip on one or more of the 30 days preceding the survey.)							
NA	NA	6.3 (±1.0)	5.1 (±1.4)	4.2 (±1.8)	5.0 (±1.2)	No change, 1995 – 2001	No change
Current cigar use (Smoked cigars, cigarillos, or little cigars on one or more of the 30 days preceding the survey.)							
NA	NA	NA	22.0 (±2.1)	17.7 (±1.7)	15.2 (±1.2)	Decreased, 1997 – 2001	Decreased
Current tobacco use (Smoked cigarettes or cigars or used chewing tobacco, snuff, or dip on one or more of the 30 days preceding the survey.)							
NA	NA	NA	43.4 (±2.4)	40.2 (±2.7)	33.9 (±2.1)	Decreased, 1997 – 2001	Decreased

¹ Based on linear and quadratic trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade.

² Based on T-test analyses.

³ Not available.

⁴ 95% confidence interval.

Where can I find more information?

More information about the YRBSS is available at www.cdc.gov/yrbss or call (888)231-6405.



DEPARTMENT OF HEALTH AND HUMAN SERVICES
CENTERS FOR DISEASE CONTROL AND PREVENTION